


UPDATED August 06, 2020

# Protected TRANSPORT

Intubated + Suspected/Confirmed High Consequence Pathogen

## WITH Patient



Transport Staff in PPE. Change gown and gloves prior to transport. Maintain mask and faceshield.

## AHEAD of Patient



Safety Lead  
(No PPE)



Safety Leader has no contact with patient or Transport Staff.

EXPERIENCED STAFF ONLY


### Required Droplet/Contact PPE (use donning/doffing checklist):

1. Level 2/yellow gown
2. Surgical mask
3. Goggles
4. Face Shield
5. Gloves




### Transfer on closed circuit. Change gown/gloves. Wipe stretcher.

All staff to keep mask and face shield and don new gloves and gown for transport. Transport does not require N95. Routine droplet/contact PPE apply (unless N95 already donned for AGMP). Disconnect non-essential equipment. Patient Transport to wipe bedrails/head board prior to transport.



### Have a clear Transport PLAN. Call Receiving Unit.

HUDDLE-UP. Notify receiving unit. Confirm equipment needed for transport.



### Safety Leader to open doors/elevators. Take SEPARATE elevators.

Safety Leader to follow during transport and will be responsible to open doors/elevators while maintaining no contact with patient or transport staff. Take separate elevator or stairs. Maintain 2m distance from team/patient at all times.



### Supervised removal of PPE.

Staff to individually, slowly and methodically doff PPE while observed by Safety Leader as per doffing guidelines and report any breaches of PPE immediately.

Review full protocols on <https://sunnynet.ca/coronavirus>

Updated 2020Aug06